

PARANEOPLASTIC SYNDROMES IN THE DOG AND CAT

Maurice Zandvliet, DVM, MVR, PhD, DIPL ECVIM-CA [Med & Onc]
UU Animal Cancer Center - Medical & Radiation oncology
faculty veterinary medicine, Utrecht university

1

PARANEOPLASTIC SYNDROMES

- All signs not related to the physical presence of the tumor mass, These indirect effects result from the production and release of biologically active substances including
 - Hormones
 - Growth factors
 - Cytokines
- PNS are:
 - Unrelated to size or location of the primary tumor
 - Unrelated to presence/absence of metastases
 - Maybe present before the actual tumor mass is noted
 - May serve as a tumor marker

The chance that a cancer-patient is brought to the clinic for PNS rather than the actual tumor is high

2

PARANEOPLASTIC SYNDROMES

<u>Constitutional</u> <ul style="list-style-type: none">• Weight loss/Cachexia• Fever• Hyperviscosity• Hypertrophic osteopathy	<u>Secondary auto-immune/immune-mediated disease</u>
<u>Cutaneous</u> <ul style="list-style-type: none">• Nodular dermatofibrosis• Superficial necrolytic dermatitis• Pemphigus vulgaris• Alopecia	<u>Hematologic</u> <ul style="list-style-type: none">• Anaemia• Polycythemia• Leukocytosis (neutrophilia; eosinophilia)• Thrombocytosis• Pancytopenia
<u>Neurologic</u> <ul style="list-style-type: none">• Myasthenia gravis• Neuropathy	<u>Coagulation disturbances</u> <ul style="list-style-type: none">• Thrombocytopenia• Thrombocytopathy• DIC
<u>Gastrointestinal</u> <ul style="list-style-type: none">• GI-ulceration• Diarrhea	<u>Metabolic</u> <ul style="list-style-type: none">• Hypercalcemia• Hypoglycemia• Hyperhistaminemia• Hypokalemia

3

WHAT IS THE PNS MOST COMMONLY ASSOCIATED WITH CANCER BY OWNERS?

4

DUSHI – DSH FN 15Y

Has received multiple surgeries for mammary gland masses (nodulectomies) with no histopathology performed


Owner recently noted:

- Weight loss
- Progressive loss of appetite
- Otherwise fine




5


THORACIC RADIOGRAPHS



6




8




9

- # CANCER CACHEXIA
1. Altered carbohydrate metabolism
 - Anaerobic glycolysis of glucose in tumor cells
 - Relative insulin resistance (higher insulin levels)
 2. Protein breakdown higher than protein synthesis
 - Muscle wasting, decreased immune and GI-function, poor wound healing
 - PIF (Proteolysis Inducing Factor)
 3. Increased lipolysis and decreased lipogenesis
 - Due to changes in lipoprotein lipase and/or peripheral insulin resistance
 - Tumors have limited ability to use lipids for energy

10



11



12

CANCER CACHEXIA - TREATMENT

Multimodal approach essential in order to achieve maximum effect!

1. Treat the cancer (if possible)
2. Reduce adverse events anti-cancer therapies
3. Reduce inflammatory response
4. Dietary interventions
 - Optimize diet
 - Supplements
 - Additional feeding
 - Assisted feeding / Feeding tubes

13


CANCER CACHEXIA - DIET

Theoretical "ideal" dietary requirements:

- Highly palatable
- Highly digestible
- High in energy content
- High In Fat
- Low in Carbohydrates (Ringer's lactate myth)
- Low in Proteins
- Additional omega-3 fatty acids (EPA/DHA)?

14

CANCER CACHEXIA



Dogs with multicentric lymphoma treated with doxorubicin and "ideal" diet:

1. Beneficial effect on DFI and survival
2. More rapid decline circulating cytokine levels
3. Reduced insulin resistance & hyperlactatemia

median: 167 days; 75th quantile: 209 days; 25th quantile: 63 days). Dogs fed the experimental diet had numerically (**P = 0.23**) greater mean $\pm$ SE (318.6 plasma; 57.0 days; median (319 days); 75th quantile ST (440 days), and 25th quantile ST (124 days) compared with the dogs fed the control diet (mean $\pm$ SE: 227.6 $\pm$ 42.1 days; median: 232 days; 75th quantile: 388 days; 25th quantile: 94 days) (Table 4). Stratifica-

Ogilvie et al, Cancer, 2000.
Effect of fish oil for dogs with lymphoma.

15

WHAT IS THE MOST COMMON PNS ASSOCIATED WITH CANCER?

16

ANEMIA

- Variable degree of anemia, typically mild to moderate
- Often multifactorial
- Signs may vary depending on degree of anemia and rate of onset

Anemia	
<u>Regenerative</u> <ul style="list-style-type: none">• Blood loss• Hemolysis<ul style="list-style-type: none">• Immune-mediated• Non-immune-mediated<ul style="list-style-type: none">• Infectious• Hypophosphatemia• Heinz body anemia• Hemophagocytic syndrome• Microangiopathic hemolysis	<u>Non-regenerative</u> <ul style="list-style-type: none">• Acute blood loss• Anemia of chronic disease• Endocrine dysfunction (EPO, T4, cortisol)• Deficiency: (relative) iron, protein, vitamins• Immune-mediated destruction of erythroid precursors (e.g. PRCA)• Myelophthisis• FeLV


17

ANEMIA IN CANCER PATIENTS

- Anemia of chronic disease**
 - typically mild, normocytic-normochromic, non-regenerative
 - relative iron-deficiency, (hepcidin) reduced life-span erythrocytes
- Myelophthisis**
 - non-regenerative
 - often bi-/pancytopenia
- Blood loss anemia**
 - microcytic, hypochromic, hypo-/non-regenerative
 - mast cell tumors, intestinal tumors, gastrinomas,
- Immune-mediated hemolytic anemia** (spherocytes)
 - Regenerative
 - haematopoietic tumors (malignant lymphoma)
- Microangiopathic hemolytic anemia** (schistocytosis)
 - Hemangiosarcomas, DIC

18


ANEMIA



Spherocytes, anisocytosis, polychromasia, nucleated red blood cells
Immune-mediated hemolytic anemia

19


ANEMIA



Secondary infections
incl. *Mycoplasma* spp
hemofelis

20


ANEMIA



Hypochromasia, low MCV, thrombocytosis, low plasma proteins
Chronic gastro-intestinal blood loss (iron deficiency)

21



ANEMIA



Bone marrow -
Phagocytosis
erythroid precursors

22

ANEMIA

Schistocytes, acanthocytes, thrombocytopenia,
nucleated red blood cells
Microangiopathic hemolysis (splenic hemangiosarcoma)

23


WHAT IS THE BEST KNOWN PNS ASSOCIATED WITH CANCER?

24

YULA - SWISS WHITE SHEPHERD, F, 8Y6M

- Presenting signs: Itching and licking around tail/anus.
- Referring vet:
 - Anal sac problem?
 - PE: small mass (<1cm) in left anal sac
 - Cytology: consistent with apocrine anal sac adenocarcinoma
- Surgery: marginal excision L-sided anal sac carcinoma
- Referred for post-operative radiotherapy
 - Rectal examination: no palpable mass
 - US abdomen and thoracic radiographs: unremarkable
 - Total calcium: 3.76 mmol/L (Ref: 1.98-2.97 mmol/L)
- So what's next?


25



26

- # HYPERCALCEMIA
- Laboratory error**
 - Non-neoplastic causes**
 - Renal failure (renal secondary hyperparathyroidism)
 - Dietary (nutritional secondary hyperparathyroidism)
 - Hypervitaminosis D (rodenticides, plants, dietary oversupplementation)
 - Addisonian's disease / hypoadrenocorticism
 - Granulomatous inflammation (vitamin D analogues)
 - Idiopathic (cat)
 - Neoplastic causes**
 - Primary hyperparathyroidism (PTH)
 - Parathyroid hyperplasia/adenoma/carcinoma
 - Humoral Hypercalcemia Malignancy (PTH-rp)
 - T-cell lymphoma, anal sac carcinoma, mammary carcinoma, hepatocellular carcinoma, other carcinomas, thymoma, histiocytic sarcoma
 - Osteoclast Activating Factors (OAF's: IL-1, TNF α)
 - Multiple myeloma, B cell lymphoma
 - Osteolytic tumors (primary or secondary)
 - Incl. osteosarcoma, feline squamous cell carcinoma, bone metastases

27



28

PTH-RP

Produced in many primitive neuroendocrine tissues and placenta

Wide variety of activities due to processing of precursor into multiple hormones and the use of multiple receptors

Endocrine (3), paracrine (mostly) and autocrine function

29

PTH-RP

PTHrP is a protein polyhormone:

- N-terminal region: 70% homology with PTH; binds and stimulates PTH receptors in bone/kidney, to lower extent 1,25-dihydroxyvitamin D production
- Midregion: stimulation Ca-uptake placenta
- C-terminal region: can inhibit osteoclastic bone resorption

Human PTHrP N-terminal RIA can be used in the dog and cat

30


HYPERCALCEMIA DIAGNOSIS

- Start with
 - Blood: total Ca (if available iCa²⁺), PO₄, albumin, kidney function
 - Urine: s.g.
 - Repeat calcium!!
- Look for cancer
 - Physical examination.: lymph nodes (incl. cytology), rectal exam
 - Imaging: thoracic radiographs; abdominal US ± parathyroids
 - Bone marrow
- May need to consider
 - PTH/PTH-rp/VitD₃

31

YULA

	Result	Reference Range
Calcium	4.56 mmol/L	1.98-2.97
Albumin	34 g/L	26-37 g/L
Urea	5.1 mmol/L	3.0-12.5
Creatinin	99 umol/L	50-129
PTH	133 pg/mL	
PTH-rp	< 0.1 pmol/L	




32

HYPERCALCEMIA - TREATMENT

- Identify cause!!!!
- Correct dehydration and start diuresis
 - Calcium-free fluids - 0.9% NaCl preferred
- Diuretics
 - Furosemide; rarely necessary
 - only after correction hypovolemia!
- Biphosphonates (Pamidronate)
- Salmon calcitonin
- Mithramycin
- Glucocorticoids

Steroids should only be used **after** a diagnosis has been made or all diagnostic samples have been secured!



33

PARANEOPLASTIC CUTANEOUS SIGNS



42


PARANEOPLASTIC - CUTANEOUS



Symmetric alopecia due to hyperestrogenism
Sertoli cell tumor

43



PARANEOPLASTIC - CUTANEOUS



Nodular dermatofibrosis
Bilateral renal cystadenocarcinomas
Genetic disorder in German shepherd dog

44

PARANEOPLASTIC - CUTANEOUS



Mucocutaneous eruptions with blisters/erosions
Paraneoplastic pemphigus

45

PARANEOPLASTIC - CUTANEOUS



Superficial necrolytic dermatitis*
Glucagonoma

- Rare condition, usually older dogs
- Progressive skin lesions (weeks to months)
- Pathogenesis unclear
 - Associated with glucagonomas, chronic hepatic disease, DM
- Low plasma amino-acid concentrations (30% of normal)
 - Humans: low zinc and essential fatty acids

*a.k.a Necrolytic Migratory Erythema or hepatocutaneous syndrome

46

PARANEOPLASTIC – CUTANEOUS SUPERFICIAL NECROLYTIC DERMATITIS

Signs

- Cutaneous lesions:
 - erythema, crusting, ulceration, alopecia
 - marked crusting and ulceration of footpads
- Facial area/oral cavity, pressure points, perianal region
- Secondary infections common

Diagnosis

- Histologic skin biopsy

Therapy

- Treat primary disease
- Supportive treatment
 - Antibiotics
 - Dietary protein supplementation with egg-yolk or high-quality protein
 - Amino acids infusions

48

DUNCAN – DSH MN 7Y



Progressive non-pruritic skin lesions

Bloodwork: normal
FeLV/FIV: negative
Fungal culture: no growth
Skin scraping: no parasites

49

DUNCAN




Dermatology service

Nonpruritic scaling, diffuse erythema, and alopecia, mostly on the head and pinnas in an otherwise healthy cat

50

DUNCAN – DSH MN 7Y



Older cats - thymoma
Feline exfoliative dermatitis

51

FELINE EXFOLIATIVE DERMATTITIS

Signs

- Cutaneous lesions
 - generalized exfoliation, scales, and crusts
 - alopecia
 - non- to mildly pruritic
- Systemic signs: lethargy, anorexia, and weight loss for several weeks

Causes

- Thymoma
- Non-thymic neoplasms, drug administration, nutritional factors

Diagnosis

- Histologic skin biopsy: interface dermatitis, mural interface folliculitis, sebaceous adenitis

52

HUMMIE – DSH FN 11Y




Systemic signs

- lethargy
- abdominal distension

Skin

- progressive, diffuse, symmetrical alopecia
- Thin, greasy and glistening skin
- Fissured footpads

53



Associated with pancreatic or hepatic/bile duct carcinoma

Feline paraneoplastic alopecia

54

BRUCE – ROTTWEILER MN 8Y

Since 2 weeks

- Progressive lameness
- Thickened, painful distal limbs
- Loss of appetite
- Occasional cough



60

HYPERTROPHIC OSTEOPATHY

Typically (large) thoracic or abdominal mass

Signs

- (sub)acute onset (<3W)
- Lameness with (symmetrical) limb swelling
- Ocular discharge, episcleral injection, lethargy

Diagnosis

- Rx limb: periosteal reaction of distal long bones
- Mass: metastatic > primary; thorax > abdomen
- Anemia, neutrophilia elevated ALP

Treatment

- Treat underlying condition
- Neurotomy vagal nerve



61

HYPERTROPHIC OSTEOPATHY



62

HYPERTROPHIC OSTEOPATHY



Musée Fragonard de l'École vétérinaire de Maisons-Alfort (Paris, France)

63

WHAT'S YOUR DIAGNOSIS?



77